




FIG. 1: (Color online) Top: Statistical distributions of avalanche sizes S for various injection rates (clip level $C=0$). The main plot shows $P_S(S)\langle S \rangle^{\alpha/(2-\alpha)}$ vs $S/\langle S \rangle^{1/(2-\alpha)}$, and the inset $P_S(S)S^\alpha$ vs $S/\langle S \rangle^{1/(2-\alpha)}$, both computed for the value $\alpha = 1.00$ that provides the best collapses. Bottom: Corresponding collapses of the pdf's of the avalanche durations T , for the value $\tau = 1.25$.

Planet, Santucci, and Ortín Reply:

In Ref. [1] we reported that both the size and duration of the global avalanches observed during a forced imbibition process follow power law distributions with cut-offs, $P_x(x) = a_x x^{-m_x} \mathcal{G}_x(x/x_c)$. While the exponent of the power law appears robust within the quoted error bars, the cut-off of the pdf's depends on experimental control parameters such as the injection rates v .

When adimensionalising the variables, $u = x/\langle x \rangle$, we observed a collapse of the various pdf's $P_u(u)$. Thus, the exponent m_x should be found equal to one, as explained in [2]. Indeed, we observed clearly an average power law exponent $\alpha = 1.00 \pm 0.06$ for the avalanche size pdf's. However, for the avalanche duration we reported a slightly larger value within the large dispersion. This can be attributed to the poorer statistics for the avalanche duration, as we explained in [1], affecting the quality of the collapse and/or the fitting procedure. This illustrates the difficulty of extracting accurate values of the power law exponents with experimental data with a limited statistics.

In order to extract reliable exponents, the right procedure –as described in [2]– is to find the power law exponent m^* that provides the best collapse of $Y = P_x(x)\langle x \rangle^{m/(2-m)}$ as a function of $X = x/\langle x \rangle^{1/(2-m)}$. In order to quantify the quality of this collapse, first, we




FIG. 2: (Color online) Isolines of the joint distributions u' vs w' , with a fit to the crest giving the value $\gamma = 1.33 \pm 0.12$.

have fitted $Y_{fit} = AX^{-m} \exp(-X/B)$ to the set of our experimental data, varying systematically the exponent m and leaving A and B as free parameters. Then, we have computed the error $\epsilon = \sum_i [\log(Y_i) - \log(Y_{fit})]^2$ as a function of m . For various velocities (clip level $C = 0$), for the size distribution, the minimum value of ϵ (thus, the best collapse) is obtained for $\alpha = 1.00 \pm 0.15$, while for the avalanche durations we obtain $\tau = 1.25 \pm 0.25$, as shown in Fig. 1. The insets display the collapse $P_x(x)x^m$ as a function of $x/\langle x \rangle^{1/(2-m)}$, showing that the scaling function is very well approximated by a decaying exponential. Then it is not difficult to show that the joint distribution of sizes and durations can be properly analyzed using $u' = S/\langle S \rangle^{1/(2-\alpha)}$ and $w' = T/\langle T \rangle^{1/(2-\tau)}$, with the values of the exponents previously found. We show in Fig. 2 that $u' \propto w'^\gamma$, where $\gamma = 1.33 \pm 0.12$.

It is important to notice that the values obtained here are in agreement with the original ones obtained by a direct fitting of the pdf's of the raw and dimensionless data. Since the actual exponents m are close to one $P(x/\langle x \rangle) = P(x)\langle x \rangle \simeq P(x)\langle x \rangle^{\frac{m}{2-m}}$ and $x/\langle x \rangle \simeq x/\langle x \rangle^{\frac{1}{2-m}}$, and due to both the experimental noise and limited statistics, the collapses previously observed were reasonably good and nearly indistinguishable from the present ones.

Finally, Ref. [2] suggests that the problem of forced-flow imbibition might belong to the quenched Edwards-Wilkinson (QEW) or equivalently the C-DP universality class [3], a conjecture based on a possible similarity of the values of the exponents. This should be taken with some caution, however, because forced-flow imbibition is a *non-local* process [4], while the QEW interfacial equation describes a *local* interfacial dynamics. Moreover, the various values of the exponents reported here are in very good agreement with the ones obtained from phase-field simulations of a *non-local* interfacial process [5].

[1] R. Planet *et al.*, Phys. Rev. Lett. **102**, 094502 (2009).

[2] G. Pruessner, Comment LPK1047 submitted to Phys.

Rev. Lett., February 2010.

- [3] M. Alava and M.A. Muñoz , Phys. Rev. E, **65**, 026145, (2002).
- [4] M. Alava et al, Adv. Phys. **53**, (2), 83-175, (2004).
- [5] M. Pradas, *Interfaces in disordered media. Scaling growth, avalanche dynamics, and microfluidic fronts*. PhD Thesis, University of Barcelona (2009).